

Some New and Previous Materials on Alpine Magmatism, Tectonics, Melt and Fluid Inclusions, and Metallogeny in Eastern Iran

A. Romanko¹, N.A. Imamverdiyev², V. Prokofiev³, I. Vikentiev³, B. Rashidi⁴

A. Savichev¹, M. Heidari⁵

¹ Geological Institute, Academy of Sciences, Moscow, Russian Federation.

² Baku State University, Baku, Republic of Azerbaijan.

³ Institute of geology of ore deposits, Academy of Sciences, Moscow, Russian Federation.

⁴ Satrap Resources, Perth, Australia.

⁵ Pars Kani, geoconsulting, Tehran, Iran.

***Corresponding Author:** A. Romanko, Geological Institute, Academy of Sciences, Moscow, Russian Federation. Email: a.romanko@ya.ru, alx.romanko@gmail.com

Abstract: The Alpine tectonic-magmatic-metallogenic events in eastern Iran, Middle East and also in Lesser Caucasus - mainly Late Cenozoic (by N.A. Imamverdiyev et al.) have some common similarities. Important geological – metallogenic +- OIL / HC correlation for the Alpine time exists (metallogeny of East Iran led by outstanding regional trio: E. Romanko, A. Houshmandzadeh, and M.A.A. Nogole-Sadat). Geological northeastern (NE) zoning and “hot” tectonics due to the African superplume activity including, probably, known delamination of lithosphere during collision of mantle lithosphere ca. 13 Ma is principal here. Intraplate alkaline-subalkaline rocks of the region studied including Quaternary real carbonatites of Hanneshin, Afghanistan were derived from enriched African superplume-related mantle sources being enriched in HFSE - Nb, Ta, Zr, Y, P, Ti. Late Cenozoic High-K calc-alkaline rocks in the Lesser Caucasus could be deal with African superplume activity too despite their subduction-related rock geochemistry. Important data exist about a general meridional-close (ca. N-S) zoning of oil / hydrocarbons (HC), muds, salts etc. here. This is one of arguments in favor of deep HC input alongside to traditional HC interpretation too. Large regional economic Cu-Au porphyry etc. metallogeny deals mainly with Eocene (Pg2) shoshonite – latite series rocks formed during subduction of Arabian plate beneath the Central Iran.

Keywords: eastern Iran, Middle East, conjunction in Alpine-Himalayan mobile belt, geology, geochemistry and petrology, tectonics, magmatism, metallogeny, African superPlume, delamination, mineralogy, melt and fluid inclusions, northeastern (NE) tectonic-magmatic-metallogenic +- oil / hydrocarbons (HC) zoning.

1. INTRODUCTION

The studied area is a small central fragment in the huge Alpine-Himalayan orogenic (collisional or mobile) superimport belt (Stocklin et al., 1965; Milanovsky, Koronovsky, 1973; Nogole-Sadat et al., 1985; Houshmandzadeh et al., 1986; E. Romanko et al, 1984; Khain, Leonov, 1988, Leonov et al, 2010; Trifonov, Ruzhentsev, 1984) and many other works. As we know, this fantastic region is very important. Geo-data in some structures of the region are poor. The first field materials were received by our team led by outstanding regional geo-trio as A. Houshmandzadeh, M.A.A. Nogole-Sadat and E. Romanko in fairly non-simple field conditions. Here are some new and previous data.

2. OUTLINE OF GEOLOGY

General geology and tectonics were characterized in such works as follows: (Nogole-Sadat, 1985; Houshmandzadeh et al., 1986; etc. and etc., fig. 1). Two groups of magmatic rocks were revealed here by our team as: mainly Eocene shoshonitic-latitic etc. rocks of the first group (subduction-related one) and principally other rocks - Neogene – Quaternary intraplate subalkaline and alkaline ones, second group (Romanko et al., 2005).

Differentiated rocks of the **first group** are the products of a large mainly Paleogene subduction of the Arabian plate beneath the Central Iran block (Fig. 1). This subduction is confirmed by the regional tectonic analysis (Leonov et al., 2010), High-resolution tomography by known J. Ritsema's team (Bull et al., 2009 etc.), geochemistry (Romanko et al., 2013; etc., fig. 1) etc. Catastrophic earthquakes up to 8M and more, unfortunately, are not rare here. A recent catastrophic precedent is 2003 Bam earthquake in East Iran with a lot of casualties. Strong seismic hazards in Pakistan, Afghanistan, Turkey etc. including the very 2015 year bring civilian and economic damage.

Antipodes of the **second group** related to African superplume activity are: intraplate K-Na subalkaline and alkaline rocks – High-Ti trachybasalts, trachyandesites, real Quaternary carbonatites of Hanneshin, Afghanistan, Late Cenozoic carbonatites of Arabia, also Neogene lamproites of Algeria etc. by E. Romanko et al., 1988 and Romanko et al., 2013 (tables 1-11, fig. 1-2, 9-11; Bogatikov et al., 1987; Luchitsky, 1985; Shilov, 1997; Yarmolyuk et al., 2001; Knipper et al., 1992 etc.)




Fig1. Volcanic associations in the region studied, with special reference to the Lesser Caucasus using data by Dilek, Imamverdiyev, and Altunkaynak, 2010, with small change and simplification. Explanation (as in the very figure): 1 – red color – post-collisional differentiated volcanites including trachybasalts etc., 2 – yellow color – mainly basalts in peri-Arabian region, 3 – continental blocks with Afro-Arabian affinity, 4 – plate convergence vectors with respect to fixed Eurasia.




Fig1-2. Distribution of earthquake epicenters - red circles - in the Middle East after (Alinaghi et al., 2007). We can see known Lut Block and immediately next to the east - studied East Iran zone and adjacent structures marked by oval.




Fig1-3. Magmatic samples position in East Iran using Geological map scale 5: 000 000. R-70, R-71 – intraplate rocks in Sistan, R-75ia - High K-dacite with a high crystallization temperature as shown in text, Carb = carbonatites of Hanneshin, Afghanistan, R-28 – Lar alkaline intrusion with Cu-Au porphyry mineralization, Abbas Abad – important area with Cu deposits, tab = basic trachyandesite, tb = trachybasalt.




Fig2-1. Na_2O+K_2O (wt %) versus SiO_2 (wt %) or TAS diagram. Triangles - intraplate rocks of East Iran, N? age, quadrangle – Lar alkaline intrusive massif with Cu-Au mineralization, Pg3? age, E. Iran/ W. Afghanistan border. Dot – trachydacite of shoshonite – latite series, Kurama zone, Tien-Shan, C3-P1, analogue of Pg2 shoshonite – latite series (Lesser Caucasus - Central Iran - East Iran).




Fig2-2. Na_2O+K_2O (wt %) versus SiO_2 (wt %) or TAS diagram for the Abbas Abad Cu-mining area, Central Iran (NE Iran by a formal geography), Pg2? age. Samples of M. Heidari et al.

These intraplate rocks, in contrast to subduction-related calc-alkaline and other rocks, are characterized by an enrichment in both LILE - K, Rb, Ba, Cs and HFSE - Nb, Y, Ta, Zr, Ti, P, etc. (Tables or tab. 1, 5-11, fig. 3) with a characteristic positive Eu/Eu^* - more than 1.0-1.1. Also, increased content of P_2O_5 - sometimes more than 1.0% (very high) - is a characteristic feature of intraplate rocks.

We have received fair low isotopic data $^{87}Sr/^{86}Sr$ (ISr) in two samples of intraplate rocks of the second type - trachyandesites R70-2 - $0.7039 \div 0,2$ (high $K/Rb=393$) and trachybasalt R71-4 - $0.70489 \div 0,18$ ($K/Rb=375$, fig. 4). For subduction-related calc-alkaline andesite of stratovolcano Bazman, sample R-25 was determined a rather low value $ISr = 0.70456 \div 0.05$, $K/Rb=250$ (tab. 1). Isotopic data of these our intraplate rocks differ from collisional and subduction-related rocks from Anatolia, Turkey (Khain, 2001; Imamverdiyev, 2008 etc.). Igneous rocks of the volcanic rocks are fully differentiated series of the regional known Sahand - Bazman belt. Known mainly andesite stratovolcanoes in this belt are: Bazman with a height 3490m and Taftan - 3940m (old mark was 4042m). Old 0.7049 isotopic date for a 'volcanite' of an unnamed volcano in a desert was reported by Canadian team (Camp, Griffis, 1982).




Fig3. Thin section in XPL. R-70 sample. Carbonate-bearing volcanite in Zabol area associated with real carbonatite in Hanne shin structure, Afghanistan. Carbonate is in individual areas of thin section, left and upper right. Up to 33.0 wt.% of CaO in same R-70 group volcanites.




Fig4. Thin section in PPL. R-82 sample, Quaternary? calc-alkaline rhyolite. Bazman stratoVolcano (N-Q age). Quartz - down of thin section. Dark grains - ore minerals.

Table1. Major- and trace-element composition in the rocks studied

Sample	1	2	3	4	5	6	7	8	9
SiO ₂	48.17	57.80	54.50	54.00	60.69	65.39	65.10	85.00	58.67
TiO ₂	2.20	1.31	1.87	1.52	0.36	0.42	0.51	0.60	1.70
Al ₂ O ₃	3.80	17.48	15.94	-	15.32	13.71	15.54	4.00	15.13
Fe ₂ O ₃	9.32	4.37	6.39	6.25	2.70	3.25	2.42	3.21	6.69
FeO	2.56	1.07	0.40	-	2.07	-	2.32	1.10	2.19
MnO	0.14	0.09	0.09	0.08	0.09	0.057	0.13	0.02	0.09
MgO	5.75	2.27	3.37	-	3.65	1.39	1.72	0.52	2.28
CaO	8.98	7.10	7.58	7.40	3.90	2.08	2.80	0.29	1.77
Na ₂ O	4.93	5.11	5.81	-	3.64	2.87	3.36	0.28	5.06
K ₂ O	1.31	1.42	1.73	1.09	4.38	4.51	4.59	0.21	2.05
P ₂ O ₅	0.23	0.61	1.05	-	0.31	0.11	0.20	0.09	0.30
Rb	30	19	20	15	145	117	109	7	47
Ba	375	293	-	292	1230	577	1597	390	557
Sr	1185	912	4470	950	870	232	359	440	263
Ni	86	53	58	59	50	7	13	10	44
Co	33	14	-	-	12	5	6	4	21
Cr	64	60	38	<64	50	16	18	11	72
V	220	95	-	-	81	63	54	55	107
Cu	63	65	64	77	69	15	11	17	33
Zn	113	88	113	98	32	40	57	8	82
Pb	5	20	51	5	20	27	22	20	10
Zr	283	232	339	217	96	158	246	136	219
Y	25	19.5	25	15	15	11	29	13	23
Nb	23	17	19	-	5.8	8	12	6	30
Sc	19	10.7	-	26.2	10	-	-	6.5	10
Th	3	3.65	-	4.84	12	-	16.7	1	12
U	1.2	0.99	-	1.31	1	-	4.62	3	3
La	44	32.4	-	30	18	-	34.0	15	35.2
Ce	101	68.3	-	63	32	-	64.5	28	64.2
Nd	-	31.4	-	-	-	-	27	-	25.0
Sm	-	6.00	-	-	-	-	5.6	-	5.1
Eu	-	2.11	-	-	-	-	1.3	-	1.9
Gd	-	5.08	-	-	-	-	4.1	-	4.8
Tb	-	0.78	-	-	-	-	-	-	0.9
Er	-	1.64	-	-	-	-	1.9	-	1.6
Yb	-	1.26	-	-	-	-	1.7	-	1.6
K/Rb	560	620	586	581	245	307	350	230	350

1 and 2 - trachybasalt (sample R71-4) and trachyandesite (sample R70-2) correspondently, Haji lake, Neogene (?), Afghan block, 3 - trachyandesite, Baluchestan, Iran (Camp, Griffis, 1982), 4 - trachyandesite, R75wp, Lut block, 5 - syenite, Lar intrusion with Cu-Au mineralization, Miocene(?) 6 - K-dacite, R75, Lut block, and 7 - trachyandesite, standard, Kurama Ridge Middle Tien Shan, Karamazar, Tajikistan, Late Carboniferous - Early Permian, using data and extrapolation from (Rusinov, Kovalenker, 1991; Razdolina, Moralev et al., 1993; Mamajanov, 2005; Romanko et al., 1989) 8 - leucorhyolite with many quartz grains and some ore minerals - sulfides, R-82, East Bazman volcano, Quaternary(?), 9 - trachyandesite, continental rift, standard, Proterozoic, Pechenga area, Fennoscandian or Baltic shield, by Romanko et al., 1989.

Table2. Chemistry of melt inclusions glass (wt %) in plagioclase (1, 3), host mineral (2, 4), host acid K-volcanite (5), rhyolite from Bazman stratovolcano, and plagioclase standards (7-9) due to A. Betekhtin, 1953.

Sample	SiO ₂	TiO ₂	Al ₂ O ₃	FeO	MnO	MgO	CaO	Na ₂ O	K ₂ O	P ₂ O ₅	Cl	S	Sum
1	74.77	0.19	12.94	0.58	0.08	0.12	1.52	3.88	3.93	0.26	0.00	0.01	98.28
2	58.69	0.01	24.77	0.23	0.00	0.01	6.68	7.22	0.49	0.00	0.00	0.01	98.11
3	74.48	0.15	14.53	0.53	0.04	0.10	1.69	3.02	4.10	0.00	0.01	0.01	98.66
4	58.36	0.00	24.71	0.28	0.02	0.05	7.15	6.90	0.46	0.04	0.00	0.01	97.98
5	65.39	0.42	13.71	2.93	0.06	1.39	2.08	2.87	4.51	0.11	-	-	-
6	85.00	0.60	4.50	3.98	0.02	0.52	0.29	0.28	0.21	0.09	-	-	-
7	58.16	-	26.57	-	-	-	8.35	6.92	-	-	-	-	-
8	56.05	-	28.01	-	-	-	10.1	5.89	-	-	-	-	-
9	62.43	-	23.70	-	-	-	5.03	8.84	-	-	-	-	-

1, 3 - melt inclusions glasses in plagioclase, 2, 4 - host minerals, 5 - hosted Hi-K-volcanite, sample R-75, 6 - leucorhyolite from stratovolcano Bazman, Quaternary(?), 7-9 plagioclase standards: 7 - andesite, SiO₂ = 58.16, empirical formula - Na_{0.6}Ca_{0.4}Al_{1.4}Si_{2.6}O₈, chemical formula andesite - (Na, Ca) (Si, Al)₄O₈, Webmineral.com, 8 - 9 - plagioclase theoretical composition: An₅₀ (8) and An₂₅ (9), by A. Betekhtin, Moscow, 1953.

Table3. Sum of gases by thermobarogeochemistry (cub. cm / kg)

Sample	Sum of gas (Cubic cm/kg)	Rock, age, notes
1. R26	0.933	subvolcanites and shallow intrusions, West Taftan volcano, diorites, probably Miocene
2. R38	1.022	Lar intrusion, Oligocene-Miocene
3. R61	0.401	ophiolites, Cretaceous
4. R85	0.655	ophiolites, Cretaceous
5. R35	12.942	Subvolcanites intruding CARBONATIC rocks, West Taftan stratovolcano, maximal contain, probably Oligocene-Miocene. Highest content.
6. R66	1.262	Young Cu-Zn-Pb mineralization with Au and Ag, Taftan stratovolcano, probably Quaternary

Sum of gases includes H₂, O₂, N₂, CO₂, CH₄, C₂H₆, C₃H₈, C₄H₁₀, C₅H₁₂, and C₆H₁₄. Temperature of Au mineralization is 220 – 278oC, Oligocene-Quaternary, important Lar intrusive massif with Au up to 25.4 ppm, T = 220–226oC by analyst R. Mudrogova, VNIIG GB or Nuclear geophysics Institute, Moscow region (E. Romanko et al., 2000). Maximum of gases are in subvolcanites intruding CARBONATIC rocks. Minimum of gases are in ophiolite mélange rocks.

Table4. ⁸⁷Sr/⁸⁶Sr (ISr) isotopic data from the rocks

Sample	⁸⁷ Sr/ ⁸⁶ Sr	Rock, mineral, age, notes
1.	0,7039+-0,0002	Trachyandesite, sample R-70-2, Hilmand (Afghan) block, maybe Neogene
2.	0,70489+-0,00018	trachybasalt, R71-4, lake Haji area, Hilmand (Afghan) block, maybe Neogene
3.	0,70456+-0,00005	calk-alkaline basic andesite, R-25-1, East Bazman volcano, Neogene-Quaternary
4.	0,7049	'volcanite' by Camp and Griffis, 1982, No data about age
5.	0,7047+-0,0003	biotite from trachybasalt, sample 64, Shurab - Galecha, Eocene
6.	0,7048+-0,0003	dacite, sample 166, Eocene
7.	0,7051	andesite, sample 206, Eocene
8.	0,7055	biotite from andesite, sample 203, Cheh-meh-Huri, Eocene
9.	0,7059	andesite, sample 193-A, no age data
10.	0,7051	biotite from dacite, sample 143, Gazu area, no age
11.	0,7043	granodiorite, sample 146, no age data
12.	0,7045	granodiorite, sample 151, no age data
13.	0,7051+-0.0003	biotite from granodiorite, Gazu area, Campanian
14.	0,7048+-0,0003	biotite from dacite, Shurab-Galecha, Eocene
15.	0,7056+-0.0002	plagioclase from dacite, Eocene
16.	0,7065+-0.0003	biotite from dacite, Kuh-Berg, Eocene
17.	0,7070+-0.0003	granodiorite, Sor-Kuh, Middle Jurassic
18.	0,7041+-0.0001	Late Cenozoic magma, ENd= +4.1 +- 0.2, Great Caucasus
19.	0,7040	Late Cenozoic magma, ENd= +3, Great Caucasus

1-3 - author's data, 4 - after (Camp, Griffis, 1982), 5-9 – Lut block, immediately west from East Iranian zone, after Sandwall E., Turkell N. Zor E. et al., 2003; 18-19 – Geat Caucasus, courtesy of I. Chernyshev, S. Bubnov, A. Lebedev et al., IGEM, RAS, Moscow.

Table5. Composition of rock-forming and accessory minerals

Sample	1	2	3	4	5	6	7	8	9	10	11	12	13
SiO ₂	55.45	54.11	55.20	43.79	44.65	46.25	46.87	53.86	54.54	64.49	68.94	68.61	0.018
TiO ₂	0.26	0.19	0.22	1.88	1.71	1.84	1.36	-	-	0.26	-	0.13	29.79
Al ₂ O ₃	1.51	1.52	1.50	11.37	7.88	7.69	6.40	29.51	28.22	17.82	17.20	17.85	0.02
Fe ₂ O ₃	-	-	-	-	-	-	-	-	-	-	-	-	-
FeO	11.66	15.73	12.7	13.68	13.05	13.39	15.00	-	-	2.79	1.12	0.96	62.11
MnO	0.24	0.34	0.28	0.28	0.27	0.24	0.03	-	-	0.16	-	-	0.17
MgO	28.21	27.01	28.14	15.16	14.23	15.32	13.68	-	-	0.85	0.15	0.15	1.60
CaO	1.89	1.16	1.93	10.55	10.48	10.42	11.51	8.97	9.99	1.87	0.47	0.70	0.08
Na ₂ O	0.32	-	-	2.08	1.31	1.48	1.40	5.55	5.59	9.90	7.09	7.55	-
K ₂ O	-	-	-	0.49	0.38	0.44	0.75	0.28	0.34	1.78	4.96	3.71	-
P ₂ O ₅	0.18	-	-	-	-	-	-	-	-	-	0.22	0.22	-

1-3 – Orthopyroxenes, 2-3 – standard bronzites, orthopyroxenes, 2 – from andesite and 3 – from Hb-norite, by H. Kuno, 1964; 4-7 – amphiboles; 8-9 – plagioclases; 10-12 – alkali feldspars; 13 – magnetite.

Table6. Major elements composition in the rocks

Sample	1	2	3	4	5	6	7	8	9	10	11	12	13
SiO ₂	48.17	49.0	52.76	54.50	56.95	57.80	35.10	44.26	46.10	56.7	60.69	61.79	85.00
TiO ₂	2.20	1.69	1.11	1.87	1.27	1.31	0.74	0.81	0.49	0.60	0.36	0.52	0.60
Al ₂ O ₃	13.80	14.1	17.44	15.94	16.40	17.48	13.48	12.70	10.30	11.1	15.32	17.10	4.05
Fe ₂ O ₃	9.32	9.10	3.14	6.39	5.28	4.37	7.53	4.81	5.10	4.90	2.70	1.16	2.51
FeO	2.56	- 0.11	5.40	0.40	0.46	1.07	0.73	0.87	-	- 0.10	2.07	3.53	1.21
MnO	0.14	9.23	0.13	0.09	0.08	0.09	0.16	0.12	0.08	4.85	0.09	0.10	0.02
MgO	5.73	7.72	5.55	3.37	3.35	2.27	5.46	6.60	9.00	12.0	3.65	3.04	0.37
CaO	8.98	3.06	8.62	7.58	6.80	7.10	26.66	17.10	15.86	1.84	3.90	5.25	1.55
Na ₂ O	4.93	1.84	3.46	5.81	5.33	5.11	0.80	2.96	0.86	1.95	3.64	4.11	0.28
K ₂ O	1.31	0.40	1.31	1.73	1.50	1.42	0.10	0.42	2.36	0.12	4.38	1.58	0.21
P ₂ O ₅	1.11		0.40	0.51	0.59	1.05	0.16	0.38	0.12		0.31	0.19	0.03

1-10 – Hilmand (Afghan) block: 1-3 – trachybasalts, 11 – syenite, Lar massif, 12 – 13 – Bazman volcano, Neogene – Quaternary, author's data; 2, 7, 10 – data by A. Houshmandzadeh and M.A.A. Nogol Sadat et al., 3 and 4 – (Camp, Griffis, 1982), '-' not determined.

Table7. Rare Earth Elements (REE) in the rocks studied and standards

Sample	1	2	3	4	5	6	7	8	9	10
La	32.4	32.1	44.8	18.6	35.2	34	63	78	31.3	23
Ce	68.3	69.3	91.9	37.7	64.2	71	115	50	50.8	43
Pr	8.23	8.05	9.80	4.32	-	-	-	-	-	-
Nd	31.4	32.9	37.8	17.7	25.0	43	70	63	21.3	
Sm	6.00	5.98	7.24	3.92	5.1	10	17	12	4.09	4.72
Eu	2.11	1.83	1.31	1.23	1.9	3.0	4.5	4.0	1.26	1.56
Gd	5.08	5.55	6.19	4.20	4.8	7.5	11	10	3.42	
Tb	0.78	0.71	0.70	0.54	-	-	-	-	0.55	1.93
Dy	3.20	3.13	3.76	3.50	-	-	-	-	-	
Ho	0.68	0.57	0.64	0.69	-	-	-	-	-	
Er	1.26	1.40	1.93	2.21	1.6	2.8	3.7	2.9	1.79	
Tm	0.31	0.26	0.26	0.32	-	-	-	-	-	
Yb	1.26	1.10	1.74	2.23	1.6	1.8	2.4	2.8	1.94	1.97
Lu	0.34	0.23	0.25	0.34	-	-	-	-	-	

1-4 - intraplate rocks in West Baluchestan: 1-2 – trachyandesites, Neogene (r70-2 and r70-23 samples, analytics by A. Houshmandzadeh and M.A.A. Nogol Sadat support); Helmand basin, 3-4 – subalkaline rocks, Lut block (r75-1 and r75-2); 1-4 - analytics by A. Houshmandzadeh and M.A.A. Nogol Sadat support; 5-trachyandesite, standard, continental rift, Paleoproterozoic, Kuetsjarvi unit, Pechenga zone, Fennoscandian Shield by A. Romanko et al.; 6-8 – basalt and dolerite (intraplate standard rocks), continental rift, Jurassic, Karoo formation, Save-Limpopo rift, Zimbabwe, E. and A. Romanko; 9 – trachyandesite, Eocene, subduction-related setting, sample BH-13 from a well, Talmessi deposit, Central Iran, courtesy of H. Bagheri, 10- trachybasalt, sill, sample Ta-39, Eocene, Lesser Caucasus, Imamverdiyev, 2010.

Table8. Composition of glass in acid volcanite R-82, East Bazman stratovolcano, T crystallization = 690oC, content of H₂O - more than 6 wt% by different methods.

Sample	SiO ₂	TiO ₂	Al ₂ O ₃	FeO	MnO	MgO	CaO	Na ₂ O	K ₂ O	P ₂ O ₅	Cl	S	Sum	Rastr
1	72.70	0.10	10.88	0.68	0.05	0.08	0.68	2.49	3.69	0.04	0.11	0.02	91.52	12 x 12
2	72.78	0.14	11.39	0.75	0.07	0.12	0.74	2.63	3.69	0.02	0.13	0.04	92.50	12 x 12
3	72.59	0.14	11.40	0.71	0.05	0.12	0.72	2.22	3.74	0.03	0.12	0.03	91.87	20 x 20
5	71.44	0.07	11.10	0.71	0.06	0.13	0.77	2.64	3.57	0.04	0.16	0.01	90.70	12 x 12
6	71.96	0.09	11.17	0.62	0.00	0.16	0.74	2.78	3.70	0.14	0.13	0.03	91.52	12 x 12
7	72.03	0.13	11.12	0.72	0.07	0.13	0.79	2.88	3.71	0.15	0.15	0.01	91.89	12 x 12
8	72.61	0.06	11.31	0.72	0.00	0.13	0.71	2.83	3.75	0.12	0.16	0.02	92.42	12x 12

1-8 – composition of glass inclusion in Quartz of acid volcanite, rhyolite R-82, East Bazman stratovolcano, T of crystallization = 690oC, High content of H₂O = 6 wt%. There are many sulfides in a sample correlated with higher content of Cu, Zn etc. in a sample R-82. Analyses led by V. Prokofiev.

Table9. Rare, trace (ppm) and major (%) elements composition

sample	Ni	Cu	Zn	Ga	Pb	Rb	Sr	Y	Zr	Fe ₂ O _{3t}	K ₂ O	CaO	TiO ₂	Cr ₂ O ₃	MnO	Ba	La	Ce
1. 75-WP	59	77	98	14	5	15	950	15	217	6.25	1.09	7.40	1.52	<64	0.08	292	30	63
2. 71-4	77	75	113	13	6	18	1138	24	245	9.96	1.10	9.73	2.49	64	0.13	376	40	111
3. 71-42	87	63	197	14	5	14	1097	22	223	9.56	1.23	10.19	2.02	<64	0.12	375	44	101
4. 71-43	82	68	110	14	6	16	1115	23	234	9.8	1.16	9.73	2.49	64	0.13	376	40	106
5. 70-32	40	131	71	9	26	5	181	18	91	7.39	0.02	34.89	0.48	<64	0.12	40	13	14
6. 70-4	43	166	164	6	20	5	505	17	146	4.72	0.32	21.32	0.63	<64	0.09	155	20	37
7. 70-5	162	86	89	10	13	20	751	13	186	5.91	1.31	10.88	1.12	0.04	0.08	310	28	57
8. 70-6	136	65	79	11	10	22	782	18	180	5.86	1.30	10.47	1.14	0.03	0.07	304	23	55
9. 70-7	49	77	86	14	13	16	992	15	208	6.01	1.09	7.90	1.50	64	0.07	319	21	69
10. 70-8	42	77	87	13	5	13	1106	16	205	6.24	1.11	8.04	1.52	<64	0.07	334	35	64
11. 70-9	38	60	83	14	6	14	875	14	183	5.11	1.55	6.54	1.27	<64	0.07	270	30	69
12. 70-10	67	80	93	16	12	16	683	9	100	5.62	1.46	7.87	1.53	<64	0.08	318	31	68
13. 70-11	52	62	92	17	8	16	943	15	215	6.21	1.30	7.04	1.64	<64	0.08	273	30	58
14. 70-12	50	85	89	15	9	17	900	15	205	6.10	1.47	7.63	1.38	<64	0.08	324	32	68
15. 70-13	57	57	79	12	14	20	917	17	201	5.96	1.37	8.19	1.36	<64	0.08	379	35	67
16. 70-14	51	60	83	19	8	15	863	18	203	5.06	1.47	6.93	1.31	<64	0.06	292	28	64
17. 70-15	67	80	93	16	12	16	683	9	199	5.62	1.46	7.87	1.53	<64	0.08	318	31	68
18. 82-5	20	70	170	18	-	93	52	36	516	7.81	4.35	0.99	1.12	<64	0.12	781	56	104

1- 17 – intraplate rocks, Baluchestan and Sistan Province: 1- Lut block (R-75wp, sample of E. Romanko), 2-4 temporary Haji lake, north from Zabol, 5-17 – unnamed volcano in desert, 18 – important calc-alkaline rhyolite R-82, T crystallization = 690oC, H2O = 6 wt.%, east Bazman stratovolcano, Quaternary ?. “-“means below resolution concentration. XRF is by TEFA-3 techniques.

Table10. ICP-MS data (ppm) on volcanites and ore sample

2	3	4	5	6
1s59	9s66	3as1	9s15	Mr3S95
50	40	45	43	24
2,3	2,1	1,6	2,8	2,21
11,3	9,7	12,1	9,7	10,8
4097	3786	4060	3769	-
137	138	165	138	166
12	6,9	17	11	44
731	1025	786	671	-
14	13	18	12	15,3
3,6	1,8	6,5	2,4	9,25
38	284	13709	62	685
95	125	91	44	-
28	27	27	26	18
101	69	104	78	131
915	1309	814	938	1077
18	18	17	17	21
144	134	130	132	133
7,5	6,9	7,5	6,9	8,2
1,1	0,74	3,1	1,0	
2,4	10	4,1	36	
639	517	534	512	
29	29	26	30	30
59	58	54	57	59
7,3	7,2	6,7	7,0	-
29	28	27	27	29
5,4	5,3	5,1	5,2	5,3
1,6	1,6	1,6	1,6	1,47
5,3	5,0	5,0	5,0	4,7

Some New and Previous Materials on Alpine Magmatism, Tectonics, Melt and Fluid Inclusions, and Metallogeny in Eastern Iran

0,71	0,67	0,67	0,67	0,69
3,9	3,7	3,6	3,6	3,88
0,75	0,74	0,72	0,71	0,77
2,2	2,2	2,1	2,1	2,21
0,32	0,31	0,30	0,31	0,32
2,2	2,2	2,0	2,1	2,10
0,33	0,32	0,30	0,30	0,34
3,6	3,4	3,2	3,3	3,35
0,47	0,42	0,45	0,43	0,83
1,8	0,49	0,82	0,73	-
16	14	14	15	10,0
0,003	0,022	0,091	0,085	-
7,6	7,3	6,9	7,2	7,9
2,4	2,6	2,4	2,0	2,03

1-6 – volcanites including Cu-rich ones, Eocene (Pg2)?, Abbas Abad

Cu mining area, NE. Iran, samples of M. Heidari.

Table11. Trace elements in rocks (ppm)

Sample	Rb	Sr	Y	Zr	Nb	Ba	V	Ni	Co	Cr	Sc	As	U	Th
1.R71-4	16	1290	25	270	38	380	240	72	35	100	25	13	<1	<1
2.R70-8	16	970	14	220	15	380	95	32	14	53	14	8.9	<1	1.8
3.R70-28	17	970	16	220	16	340	100	36	13	54	13	10	<1	2.9
4.R70-27	3.8	510	12	130	11	140	89	26	12	72	16	19	<1	3.3
5.70.272	4.1	170	16	75	5.5	120	130	20	2.8	21	27	17	4.2	3.6
6.R28-59	145	1230	13	110	5.8	870	81	50	12	55	15	7.9	6.3	12
7.34-24	6.7	170	23	85	8.4	76	34	7	5.3	39	5	4.8	1	14

R-71-4 - trachybasalt, samples R-70 – trachyandesites and associated intraplate rocks, 28-59- syenite, Lar, intrusive massif, NI?, 34-24 – acid subduction-related dacite, N. Pamirs, Late Permian (P2), V. Novikov, A. Romanko et al, 1992, for comparing. XRF, ppm, Geological Institute, RAS.




Fig5. The distribution of the contents of rare and trace elements normalized to chondrite composition (Sun, McDonough, 1989), using (Saadat S, Stern C.R., 2011).




Fig6. Isotope systematics of igneous rocks in the region and standards using (Saadat S, Stern C.R., 2011).

3. INCLUSIONS

Melt inclusions in this region were firstly investigated under the leadership of Prof. Prokofiev, IGEM RAS as well as fluid inclusions by E. Romanko et al. in 2000. Some notes and conclusions here are as follows:

- Melt inclusions are not typical for the African super-plume-related intraplate igneous rocks. Intraplate rocks are confirmed by a tomography of known Ritsema's team (Bull et al., 2009 etc). Also, melt inclusions are not typical or rare for shoshonite series rocks of Abbas Abad area. T crystallization of melt inclusions in similar Eocene shoshonite series rocks with Fe-skarn mineralization, West Iran is fairly high - ca 300oC by V. Prokofiev et al.

- unusual fairly high temperature, 1150-1180° C - up to 1220° C melt inclusions were revealed in plagioclase of subduction-related K-dacite, sample 75-1 by V. Prokofiev et al, 2011 (Prokofiev, 2000; Romanko et al., 2012, Fig. 5 and 6, Tables 2, 8.). This fairly deep, non-calc-alkaline rock was also affected by indirect (?) influence of a huge African super-plume, as proposed. Homogenization occurs under High T = 1150-1220° C (for comparing, for example, T much lower for acid volcanite of Quaternary Pektusan volcano, Korea, paper of O. Andreeva et al., IGEM RAS, Moscow, 2013). A higher viscosity of a glass provides more inclusions coexistence in a sample.

Maximal concentration on fluid CH₄ and other CH-based **fluid inclusions** were revealed in shallow intrusions on the contact with carbonate-rich host rocks in west Taftan zone; also in important Lar syenite massif with Cu-Au mineralization (Table 3, E. Romanko et al., 2000). Opposite, minimal data are in Cretaceous ophiolitic mainly melange rocks.



Fig7. Sample N-40. L. Caucasus, for comparison. T of crystallization = 1080-1140°C. Melt inclusions in Plagioclase (Plg) from trachyandesite. Similar melt inclusions are in similar Alpine volcanites in eastern Iran too.




Fig8. Sample R-75ia. Baluchestan and Sistan Province, Iran. $T=1150^{\circ}\text{C}$. View of melt inclusions in acid glass from Plagioclase




Fig9. Sample R-75ia. T of crystallization is fairly high - 1150-1220oC. Homogenization of melt inclusions.




Fig10. Sample R-82. Melt inclusions in Quartz, acid rhyolite, Bazman strato-volcano, $T = 690^{\circ}\text{C}$. H_2O content is high = 6 wt%. Naturally chilled melt inclusion. Maximal size of inclusion is ca 60 μm or microns.



Fig11. Sample R-82. Similar melt inclusion with gas bubbles in Quartz, rhyolite, East Bazman volcano, $T = 630^{\circ}\text{C}$. After next $T = 690^{\circ}\text{C}$ gas bubbles will disappear. H_2O content is up to 6 wt%.




Fig.12. Spider-diagram for non-intraplate rocks, principally other, probably - subduction-related rocks. Circles – shoshonite series rocks, Abbas Abad copper deposit, Central Iran, M. Heidari's samples, cross – intraplate rocks of Lut block by Saadat et al, 2002, dots – Kurama Mt, Tien Shan, C3-P1 age, analogues of studied shoshonitic – latitic rocks. Usual positive anomaly by Pb. Ta – Nb deficit here is in subduction-related? rocks. For a comparison, example: Cross – principally OTHER intraplate rocks of Lut block by Saadat et al, 2002.




Fig13. Spider-diagram for intraplate rocks of East Iran, typical flat profiles. Sometimes -usual Pb positive anomaly. Triangulars - R-70 – unnamed volcano (full plot with usual Pb – positive anomaly), R-71 – Haji temporary lake in a desert, not-full profile, no Middle - HREE data here, cross – intraplate rocks of Lut block by Saadat et al, 2002.




Fig14. Chondrite-normalized REE diagram for Cenozoic intraplate rocks and Eocene (Pg2) subduction-related ones of Iran, N=6. Intraplate rocks are: R-70 (triangulars), R-71 (filled green triangulars), and crosses - sample from Lut block (Saadat et al., 2002). Circles – shoshonite – latite series rocks, Eocene (Pg2), Abbas Abad Cu mining area, N=2, samples of M. Heidari. Dots – rock of shoshonite- latite series, analogue, Kurama Mts, Middle Tien Shan, Late Carboniferous – Early Permian (C3-P1). Absent of Eu-deficit is typical for both intraplate and subduction-related rocks.

Intraplate rocks were derived from deeper mantle source versus subduction-related Eocene and Late Cenozoic rocks. This is supported by the following:

- Geological and petrographic and mineralogical data
- The general style of petrology and geochemistry of these rocks, we can the same on other regions
- Obvious geochemical materials, for example, the stable high $K/Rb = 560-586-620$ etc.

The region is expected to at least partial compensation of Pg2-Q aged compression and subduction-related Magmatism by intraplate magmatism. The latter, according to the imaging may be associated with the tail of the most powerful African superplume (Bull et al., 2009). There is also discussion in modeling - the partial screening of the plume push up plate, which is not an obstacle - it is known that plate moves may not stop movement of the tail superplume in lateral direction.

4. METALLOGENIC NOTES

Neogene rocks of the Lesser Caucasus is interesting for economic and non-economic metallogeny as:

- New Low-temperature Au-Ag, Hg, As, Sb, Cu-Mo with Au, Cu-Pb-Zn and Pb-Zn deposits and occurrences is proposed here,
- Non-ore raw – tuffs, slags, pumice etc. are of interest too.

Metallogeny of Cenozoic rock of East Iran was studied under the leadership of outstanding regional trio – Dr. Eugene Romanko, Dr. A. Houshmandzadeh, and Dr. M.A.A. Nogole-Sadat.

Calc-alkaline intrusive, extrusive, pyroclastic and volcanogenic-sedimentary rocks here are characterized by a common copper-gold (Cu-Au) metallogenic profile for Baluchestan and Sistan Province in East Iran, as in the whole Sahand – Bazman volcanic-plutonic belt of Iran. The overwhelming majority of occurrences the study area is associated with magmatic complexes. Such metallogenic types were revealed here as:

- Multi-sulfide (Au-Mo-Cu-Pb-Zn) subvolcanic porphyry type;
- Au-As-Hg-W-Mo-volcanic exhalation one;
- Low-sulfide gold-silver plutonic one;
- Gold-copper (Au-Cu) skarn and plutonic-hydrothermal one (E. Romanko et al., 2000; data by Pars Kani Co, 2003 by Daliran et al., 2005) using also known data on mineralization of different region including former USSR / CIS (Prokofiev et al., 2000; Vikentiev et al., 2004 etc.);
- Sulfide, sulfur, alunite exhalation, surface one;
- Native-copper-sulphide volcanogenic one with zeolites;
- Silver volcanogenic sulphide (+ gold?) one.

Thus, intraplate rocks are strongly specialized in REE, P (usual process), then in Sr, Ba, U, Th due to nowadays materials. So, tectonic-magmatic, and as revealed E. Romanko – metallogenic zonation in the region was revealed in the region studied (at least in the Central – East Iran). Younger magmatic products are in the northeast of region due to lithosphere subduction and decreasing of African superplume activity in the same direction. Subduction-related (1 group of rocks) dominated calc-alkaline rocks and shoshonites-latites. , and, intraplate African superplume-related (Laverov et al., 2004; Yarmolyuk, personal communication, 2013, etc.) midalkaline – alkaline rocks including known Pleistocene carbonatites of Hanneshin, Afghanistan and, also, of one of Arabia are subordinated (2 group of rocks). Rocks of 1 and 2 groups are interpreted by us as a tectonic-magmatic couple due to one from physics etc. In this case, at least, partial compensation of subduction compression by the intraplate extension is possible. Cenozoic intraplate rocks intraplate carbonatite-derived depth of the melt - an argument in favor of the African superplume influence on the magma plume of a large region, which is in agreement with effective tomography of the well-known J. Ritsema's team (Bull et al., 2009).

5. OIL AND GAS, HYDROCARBONS (HC), SOME NOTES

There are known materials about of Caspian Sea OIL / HC resources or productivity decreasing in north – northeast (N - NE) tectonic direction or lineament as stressed by known Prof. V. Khain with

co-authors in the Explanatory map of Caspian Sea region scale 1:2 500 000 etc. (Khain, 2001; Leonov et al., 2010). This decreasing is as follows: from extremely rich Persian Gulf to South – Middle – North Caspian Sea. It is in agreement with the increasing distance from the African superplume by effective tomography (Bull et al., 2010), tectonics etc.

More specifically, this HC super belt is as Persian Gulf – Russian Arctic coast, due to old Russian HC maps, ex., USSR oil structures map scale 1:2 500 000 etc. Also, important as HC traps salt domes in the east Persian Gulf are oriented due to this tectonic direction. Relation of HC field with faults is obvious. Other possibility for HC fields is combining of well-known H₂ fluxes from the depth on faults with organic C in sediments.

Some experienced HC specialists believe that there are no strong contradictions in combined biogenic and abiogenic data now. HC fields are only of Cenozoic age, maybe the very Quaternary age for oil and younger age for the gas (not Riphean - Paleozoic – Mesozoic fields) due to a high mobility of HC. ex. There are data about ca 1 m/year HC migration. A. Timurzиеv (Timurzиеv, 2007, 2015 etc.), S. Marakushev, 2015; R. Seyful-Mulyukov, 2013 etc. think about the very deep abiogenic HC versus biogenic ones. In this context materials on mud-volcano-like structures on Mars comparing to known muds of Azerbaijan, Iran etc. by Skinner and Mazzini, 2009 etc. (fig.). It is well-known that mud-volcanoes or simply muds directly deal with HC fields.

No doubts about fault-related HC fluid input and also magmatic heat input sometimes. But biogenic factor is of great importance and **main**, surely. More data on HC peculiarities in the region studied needed.




Fig15. Comparison of Azerbaijan, Iran (L), Java mud-volcanoes (muds) with mud-like structures of Mars - after Skinner and Mazzini, 2009. Mud directly deals with OIL and other hydrocarbons.

6. CONCLUSIONS

1. Some common geo-similarities on Cenozoic events in the region studied were revealed. At least, in East Iran important north-east tectonic-magmatic zoning and partly, metallogenic one (metallogeny under the leadership of known regional trio as E. Romanko, A. Houshmandzadeh, and M.A.A. Nogole-Sadat) due to African superplume activity exists here. It caused directly by known subduction of the Arabian plate under the Central Iran. African superplume activity strongly controls magmatism, “hot” regional tectonic regime, and, at least, partly - metallogeny in the region studied. Also, African superplume closely deals with known Jurassic Karoo flood basalts

event in Africa, Paleogene magmatism in East Africa, since mainly Eocene (Pg2) subduction in Iran – Turkey etc., Neogene 11-9 Ma opening of Red Sea etc., and probably, delamination of a mantle lithosphere in East Mediterranean in Miocene and as a result – lack of regional Cu-porphphy mineralization versus economic one in Eocene before important delamination.

2. Two different types of Cenozoic magmatic rocks exist in the region studied: 1 – **intraplate** alkaline and subalkaline rocks and 2 – shoshonite - latite series rocks and calc-alkaline ones mainly dealing with **subduction – collision events**. Low crystallization temperature – 690oC and High H₂O content up to 6 wt. %, and natural melt chilling were revealed for a probably Quaternary subduction-related rhyolite of the Bazman volcano (all data on melt inclusions led by V. Prokofiev) Sudden high/very high crystallization temperature, up to 1220oC on melt inclusions in High-K probably subduction-related dacite of remnant subduction were received too. Otherwise, for intraplate rocks as well for shoshonite – latite subduction related ones melt inclusions are not typical due to proposed warm conditions.
3. Eocene (Pg2) subduction-related shoshonite – latite series rocks almost in the whole region are characterized by an economic Cu-Au mineralization with a subordinate different mineralization (Cu-Pb-Zn-Au-Ag, Hg-As, Au-Ag low-sulphide, Ag-sulfide with Au (?) etc.). Cu mineralization deals with deep basic enriched water-containing source. Cu mineralization disappears with time and higher in general magmatic section after disappears of relation with deep enriched source using (Haschke et al., 2010). Intraplate rocks bear, at least, REE, P, also Sr, Ba, Th, and U mineralization.

ACKNOWLEDGMENTS

We are extremely grateful to outstanding regional trio - E. Romanko†, A. Houshmandzadeh, and M.A.A. Nogole-Sadat† for their leadership in the field works and comprehensive care; also – to famous V.E. Khain†, M. Goncharov†, and N. Bozhko (consulting, discussion, advices); V. Trifonov, V. Burtman, I. Pospelov, M. Kopp, E. Leven and V. Volkov; Yu. Leonov†, M. Leonov, M. Fedonkin, M. Khutorskoy, B. Polyak and V. Lavrushin; V. Yarmolyuk, D. Alexeev, M. Pevzner, M. Buyakayte, D. Astafiev, M. Rodkin, I. Gablina, Yu. Malinovsky†, S. Palandzhyan and G. Nekrasov; A. Knipper†, S. Ruzhentsev†, V. Kovalenker, V. Rusinov†, A. Giris, A. Mokhov, N. Tarasov, S. Bubnov and A. Dockuchaev; G. Moralev, G. Gogonenkov et al., N. Panasuk†, also to A. Meskhi† from Kazan, Tatarstan; M. Hosseini, M. Ziaii, and more other geo-specialists for long-term discussion, consulting analytical help etc.

The research was supported by the State program no. 0135-2018-0034 (Geological Institute, Russian Academy of Sciences; N. Chamov, and partly supported by the Russian Foundation for Basic Research, project no. 17-05-0000).

REFERENCES

- [1] Abdullah S., Chmyrev V., Dronov V., Ed., 1980. Geology and Mineral Resources of Afghanistan. Moscow: Nedra. Book 1, p. 535, Book 2, 336 p. (in Russian).
- [2] Alinaghi A., Koulakov I., Thybo H., 2007. Seismic tomographic imaging of *P*- and *S*-waves velocity perturbations in the upper mantle beneath Iran. *Geophysical J. International*, 169, pp. 1089-1102.
- [3] Artemiev M., 1971. Some features of the deep structure of the Mediterranean type basins according to isostatic gravity anomalies. *Bull. Moscow Society of Naturalists, Dept. Geol.*, N 4, pp. 30-41 (in Russian).
- [4] Bogatkov O.A, Kovalenko V.I., Yarmolyuk V.V., Bubnov S.N., 1987. Series of igneous rocks: problems and solutions. *Reports of the Academy of Sciences of the USSR, Series Geol*, N 3, pp. 3-12 (in Russian).
- [5] Bortnikov N.S., Prokofyev V.Yu., 2007. Major mesothermal gold ore deposit of Russia: composition and origin of ore-forming fluids. *Digging Deeper. Proceedings of the Ninth Biennial meeting of the society for geology applied to mineral deposits*. Dublin, Ireland. 2007. Dublin Published Irish Association for Economic Geology, pp. 793-796.
- [6] Bortnikov N.S., Vikentiev I.V., 2004. Modern ore minerals formation in the oceans. *Ore deposits and ore-forming processes*. M. IGEM, pp. 325-356. In Russian.
- [7] Bull A.L., McNamara A.K., Ritsema J., 2009. Plume clusters and thermochemical piles. *Earth Planet. Sci. Lett.* Vol. 278, pp. 152–162.
- [8] Burtman V.S., 2012. Tien-Shan and High Asia. *Geological Institute RAS Transactions*. M: GEOS, 188 p. In Russian.

- [9] Dilek Y., Imamverdiyev N.A., Altunkaynak S., 2010. Geochemistry and tectonics of Cenozoic volcanism in the Lesser Caucasus (Azerbaijan) and the Peri-Arabian region: Collision-induced mantle dynamics and its magmatic fingerprint. *International Geology Review*, vol. 52, issue 4-6, pp. 536-578.
- [10] Haschke M., Ahmadian J., Murata M., McDonald I., 2010. Copper mineralization prevented by arc-root delamination during Alpine-Himalayan collision in Central Iran. *Economic Geologists*, v. 104, pp. 885-895.
- [11] Hushmand-Zadeh A., Nabavi M.H., 1986. *Metamorphic Map of Iran Scale 1: 2,500,000*. Tehran: Geol. Survey. Iran.
- [12] Imamverdiyev N.A. 2000. *Geochemistry of Late Cenozoic volcanic complexes Lesser Caucasus Baku*. Nafta-Press. 2000. 192 p.
- [13] Imamverdiyev N.A. 2008. Delamination of subducted lithospheric slab as the cause of Late Cenozoic volcanism in the Lesser Caucasus. *News of State Baku University. Natural sciences Series*. no 3, pp.536-578.
- [14] Khain V., 2001. *Tectonics of Continents and Oceans (year 2000)*. Moscow: Scientific World, 2001. 606 p. In Russian.
- [15] Khain V., Goncharov. M., 2006. Endogenic processes of different tectonic settings. *Herald of MSU*. Vol. 5. pp. 15-23. In Russian.
- [16] Knipper A., Dobretsov N., Bogdanov N., 1992. Ophiolites and "orogenic lherzolites" of Bet Cordillera, Spain. *Reports of the Academy of Sciences. Geological Series*. N 12, pp. 8-24. In Russian.
- [17] Koronovsky N.V., Demina L.I., 1999. Collisional stage of the Caucasian sector Alpine fold belt: geodynamics and Magmatism. *Geotektonika*, no 2, pp. 17-35. In Russian.
- [18] Koronovsky N.V, Lomize M.G., 2006. Tectonic accretion and collision as the steps of forming intra-continental fold belts. *Areas of active tectonics in modern and ancient history of the Earth. Materials of 39 Tectonic meeting*, vol. 1, Moscow: GEOS. pp. 353-357. In Russian.
- [19] Laverov N., Kovalenko V., Yarmolyuk V. et al., 2006. The latest volcanic activity in Northern Eurasia: Zoning and environment of formation. *Russian Academy of Sciences Reports*. vol. 410. no 4, pp. 498-502. In Russian.
- [20] Leonov Yu. G., Edit., 2007. *Alpine history of the Great Caucasus*. Moscow: GEOS. 368 p. In Russian
- [21] Leonov Yu., Volozh Yu., Antipov M. et al., 2010. Consolidated crust of Caspian region. *M. GEOS*. 64 p. In Russian.
- [22] Luchitsky I.V., 1985. *Palaeovolcanology*. Moscow: Science (Nauka), 235 p. In Russian.
- [23] Marakushev A.A., 1988. *Petrology*. Moscow. MSU publishing, 205 p. In Russian.
- [24] Milanovsky E.E., Koronovsky N.V., 1973. *Orogenic volcanism and tectonics of the Alpine belt of Eurasia*. Moscow: Nedra, 280 p. In Russian.
- [25] Nogole-Sadat, M.A., 1985. *Les Zones de Decrochement et les Virgation Structurales en Iran: Geol. Survey of Iran, Report*, 55, 259 p.
- [26] Pars Kani Co, 2003. Preliminary report on the prospection of epithermal gold mineralization in Chahnali north of Bazman, Sistan-va-Baluchestan province, Southeast Iran. *International Report*. 136 (in Farsi language with English abstract).
- [27] Pospelov I., 2009. Tectonics of Pamir. *Tectonics and geology*. V. 5, pp. 15-25.
- [28] Prokofiev V.Yu., 2000. Geochemistry of ore-forming fluids of hydrothermal gold deposits of different genetic types according to the study of fluid inclusions. Moscow. Science (Nauka), Novosibirsk, 192 p. In Russian.
- [29] Prokofiev V.Yu., Bortnikov N.S., Zorina L.D. et al., 2000. The Darasun intrusive-related gold-base metal deposit, Eastern Transbaikalia, Russia: Petrochemical, melt and fluid inclusion, REE and stable isotope (O, C, and S) studies. *Applied Mineralogy*. Rammlmair et al., Eds. Balkema, Rotterdam, Brookfield, 399-402 pp.
- [30] Romanko E., Kokorin Y., Krivyakin B. et al., 1984. *Outline of metallogeny of Anarak Area (Central Iran)*. Ministry of mines and metals. Geological Survey of Iran. Rep. 21, Tehran. Technoexport, 136 p.
- [31] Romanko E., Voinkov D., Houshmandzadeh A. et al., 2000. Characteristic features of ore mineralization in the Sistan-Baluchestan Province, SE Iran: evidence from fluid inclusions data. *International conference on geology of Mediterranean belt. Abstracts*. Beograd.
- [32] Romanko E., Houshmandzadeh A., Nogol-Sadat M.A.A. et al., 2000. Metallogenic zonation in Seistan and Baluchestan Province, East Iran. *Transactions (Doklady) of Russian Academy of Sciences*. V 370, N 1, pp. 76-79, in Russian.

- [33] Romanko E.F, Romanko A.E., Meskhi A.M., 1998. Geology and Geochemistry of Mesozoic-Cenozoic Magmatic Formations in Northeastern Algeria. Transactions of the Russian Academy of Sciences, V 362, N 7, pp. 925-927, in Russian.
- [34] Romanko A., 2005. New data on Cenozoic subalkaline intraplate rocks the East Iran. Reports of the Russian Academy of Sciences. Earth Sciences Section. V 404, N 7, pp. 510-513.
- [35] Romanko A., Imamverdiyev N.A., Prokofiev V., Vikentiev I., Savichev A., Stepanov S., 2013. Alpine tectonic-magmatic-metallogenic peculiarities in West Baluchestan, Middle East: new data, "hot" tectonics, inclusions, discussion, hydrocarbons (oil, HC), and constraints. Universal Journal of Geoscience, V 1, pp. 130-138.
- [36] Romanko A., Imamverdiyev N. A., Prokofiev V. et al. 2012. On Cretaceous – Quaternary tectonic – magmatic - metallogenic peculiarities in East Iran and South Tajikistan: new data melt inclusions, constraints, and some problems. European Seismological Commission Assembly. ESC. Book of abstracts. M. Polygraph, pp. 154-155.
- [37] Rusinov V.L., 1985. Propilites and propilitization. M.: Nauka. 185 p. In Russian
- [38] Sandwall E., Turkell N., Zor E. et al., 2003. Shear wave splitting a young continent collision. Geophys. Res. Lett., 30, no 24 b, pp.185.-194.
- [39] Shilov V.N, 1997. Comparison of Neogen-Quaternary volcanism on Sakhalin and East Sikhote-Alin. Volcanology and Seismology. V. 18. pp. 517-528.
- [40] Timurziev A.I., 2007. On a creating of new oil-gas geology paradigm based on deep-filtration model. Geophysics. N 4. pp 49-60. In Russian.
- [41] Trifonov V.G., Ruzhentsev S.V., 1984. Tectonic napes in East Pamir, Tajikistan and tectonic layering. Episodes. 32 IGC. Special Issue. pp. 15-20.
- [42] Vikentiev I.V., Yudovskaya M.A., Mokhov A.V. et al. 2004. Gold and PGE in massive sulphide ores of the Uzelginsk deposit, Southern Urals, Russia. Canadian Mineralogist. V 42, N 5, pp. 651-665.
- [43] Yarmolyuk V.V., Kovalenko V.I., 2003. Deep geodynamics, mantle plumes and their role in Central-Asian mobile belt formation. Petrology. V 11. N 6. pp. 556-586. In Russian.
- [44] Yarmolyuk V.V., Kovalenko V.I., Bogatikov O.A., 1990. South Baikal "hot spot" mantle and its role in the formation of the Baikal Rift region. Reports of Russian Academy of Sciences. V 312. N 1. pp. 187-191. In Russian.

AUTHORS' BIOGRAPHY



Alexander Romanko, Dr, Senior Researcher, Member of volcanology and chemistry of the Earth's Interior. International Association on



Nazim A, Imamverdiyev, Professor, Dr, well-known specialist in magmatism, metallogeny etc.



Vsevolod Yu, Prokofiev (Prokof 'ev), Professor, Dr, known specialist in Magmatism, metallogeny etc.



Ilya Vikentiev, Professor, Dr, known specialist in Magmatism, metallogeny, Urals, Altai etc.



Bahman RASHIDI, senior geologist, specialist in metallogeny etc.



Alexander SAVICHEV, Dr., Professor, Laboratory Chief, known specialist in chemistry, environmental sciences, magmatism, metallogeny, etc.



Mehrdad HEIDARI, senior geologist, specialist in metallogeny etc.

Citation: A. Romanko et.al. (2018). *Some New and Previous Materials on Alpine Magmatism, Tectonics, Melt and Fluid Inclusions, and Metallogeny in Eastern Iran*, *International Journal of Mining Science (IJMS)*, 4(1), pp.11-28, DOI: <http://dx.doi.org/10.20431/2454-9460.0401002>.

Copyright: © 2018 A. Romanko et al.,. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited